

Hypotonic-hyporesponsive episode (HHE)

for Health Professionals...

Hypotonic-hyporesponsive episode (HHE), is an uncommon reaction to diphtheria, tetanus and pertussis-containing vaccines (and even more rarely to other vaccines).¹ HHE occurs much less frequently following administration of acellular pertussis vaccines (aP) than with the older whole cell pertussis (wP)^{2,3} vaccines which were used in New Zealand until 2000.⁴ While an HHE may appear alarming at the time, research shows it does not result in any long term sequelae.^{3,5}

Signs and symptoms of HHE

The event is characterised by a sudden episode of limpness and unresponsiveness within 12 hours of receiving a vaccine. The average duration of the episode usually ranges from 1-30 minutes, but may last longer. For the event to be considered an HHE, all of the following must be present: limpness, unresponsiveness, pallor and cyanosis.^{4,6} Shallow breathing can also occur.⁷

Rate of occurrence

An HHE is five times more likely to occur after the older whole cell pertussis vaccine than with an acellular pertussis vaccine.⁸ Most reported cases affect infants and children under 2 years of age⁶ and usually only with the first vaccination event.^{9,10} HHE occurred 5 - 8 per 10,000^{9,11} following the older whole cell DTP vaccinations.

Sequelae

An HHE always fully resolves. No adverse outcomes have been observed after these episodes.⁶ Children who have experienced an HHE following a pertussis-containing vaccine have not demonstrated evidence of neurological damage or intellectual impairment.^{2,5} Research shows that it is very rare for a child to experience a second HHE in a child when immunisation is continued.^{6,9,12}

Follow-up care

An HHE should always be reported to the Centre for Adverse Reaction Monitoring (CARM) by the health care professional, parent or both. A documented episode constitutes a precaution (rather than a contraindication) for further doses of the vaccine involved. The affected child should be referred to their GP for assessment. A detailed record of the event should also be recorded into the child's clinical notes.

Re-vaccination

Following an HHE the immunisation programme should be completed. Having an HHE is not a reason to delay the next scheduled immunisation.

For further information contact the Immunisation Advisory Centre: 0800 IMMUNE or (0800 466 863).

To report an HHE contact CARM at: <http://carm.otago.ac.nz/index.asp?link=news&item=19>

References

1. Bonhoeffer J, Gold MS, Heijbel H, Vermeer P, Blumberg D, Braun M, et al. Hypotonic-Hyporesponsive Episode (HHE) as an adverse event following immunization: case definition and guidelines for data collection, analysis, and presentation. *Vaccine*. 2004;22(5-6):563-8.
2. Le Saux N, Barrowman NJ, Moore DL, Whiting S, Scheifele D, Halperin S. Decrease in Hospital Admissions for Febrile Seizures and Reports of Hypotonic-Hyporesponsive Episodes Presenting to Hospital Emergency Departments Since Switching to Acellular Pertussis Vaccine in Canada: A Report From IMPACT. *Pediatrics*. 2003;112(5):e348.
3. Baraff LJ, Shields WD, Beckwith L, Strome G, Marcy SM, Cherry JD, et al. Infants and Children With Convulsions and Hypotonic-Hyporesponsive Episodes Following Diphtheria-Tetanus-Pertussis Immunization: Follow-up Evaluation. *Pediatrics*. 1988;81(6):789.
4. Ministry of Health. Immunisation Handbook 2006. Wellington.
5. Martins RM, Camacho LAB, Lemos MCF, Noronha TGd, Carvalho MHCd, Greffe N, et al. Incidence of hypotonic-hyporesponsive episodes associated to the combined DTP/Hib vaccine used in Brazilian National Immunizations Program. *J Pediatr (Rio J)*. 2007;83(6):523-8.
6. Buetcher M, Heinger U, Braun M, Bonhoeffer J, Halperin S, Heijbel H, et al. Hypotonic-hyporesponsive episode (HHE) as an adverse event following immunization in early childhood: Case definition and guidelines for data collection, analysis, and presentation. *Vaccine*. 2007;25(31):5875-81.
7. Goodwin H, Nash M, Gold M, Heath TC, Burgess MA. Vaccination of children following a previous hypotonic-hyporesponsive episode. *Journal of Paediatrics & Child Health*. 1999;35(6):549-52.
8. David S, Vermeer-de Bondt PE, van der Maas NAT. Reactogenicity of infant whole cell pertussis combination vaccine compared with acellular pertussis vaccines with or without simultaneous pneumococcal vaccine in the Netherlands. *Vaccine*. 2008;26(46):5883-7.
9. Vermeer-de Bondt PEMD, van der Maas NATMD. The Effect of Age and Dose Number on the Risk of Collapse (Hypotonic-hyporesponsive episode) After Pertussis Vaccination *Pediatr Infect Dis J*. 2008;27(4):355-7.
10. Braun MM, Mootrey GT, Salive ME, Chen RT, Ellenberg SS, the VWG. Infant Immunization With Acellular Pertussis Vaccines in the United States: Assessment of the First Two Years' Data From the Vaccine Adverse Event Reporting System (VAERS). *Pediatrics*. 2000;106(4):e51.
11. Plotkin SA, Orenstein WA, Offit PA. Vaccines 5th ed. Philadelphia, Pa: Saunders; 2008.
12. DuVernoy TS, Braun MM. Hypotonic-hyporesponsive episodes reported to the Vaccine Adverse Event Reporting System (VAERS), 1996-1998. *Pediatrics*. 2000;106(4):E52.